



Study of Naturally Occurring Radioactive Materials and Some Heavy Metals from Intake of Some Nuts and Seeds in South-Western, Nigeria

Augustine Kolapo Ademola^{1*} and Ruth Omoseeke Morakinyo¹

¹*Physical Sciences Department, College of Natural and Applied Sciences, Bells University of Technology, Ota, Nigeria.*

Authors' contributions

This work was carried out in collaboration between both authors. Authors AAK conceive the idea of the study, design the methodology used, validate the data collection, write the initial draft and supervise the student project. Author ROM carried out the data collection, sample analysis, data analysis, write the revision and raise fund as student. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/PSIJ/2020/v24i630195

Editor(s):

- (1) Dr. Aleksey Anatolevich, The International Institute of Zakharenko Waves, Russia.
- (2) Dr. Olalekan David Adeniyi, Federal University of Technology, Nigeria.
- (3) Dr. Thomas F. George, University of Missouri-St. Louis, USA.

Reviewers:

- (1) Shakha Sharda, Post Graduate Government College, India.
- (2) Heba A. Saudi, Azhar University, Egypt.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/58704>

Original Research Article

Received 27 April 2020

Accepted 04 July 2020

Published 16 July 2020

ABSTRACT

The concentration of NORMs and some toxic heavy metals in some nuts and seeds commonly consumed in the south-western states of Nigeria were investigated. The analysis was done with NaI(Tl) spectrometry and Inductively Coupled Plasma (ICP) respectively. The concentration of ²²⁶Ra, ²³²Th, and ⁴⁰K in the samples was found to be below the world average. The mean values are 6.6±1.8, 3.6±1.0 and 98.2±13.5 Bq/kg, in nuts and 8.4±2.6, 2.6±1.3 and 97.6±15.0 Bq/kg in seeds respectively. The effective dose in nuts was calculated to be 13.99 and 12.0 µSv in seeds. The fatal and hereditary cancer risk estimated from the consumption of the nuts and seeds are lower than 1.0 x 10⁻⁶ which is the lowest limit. Concentration of metals in the samples descend as Fe > Zn > Cu > Cr. The Hazard Indices of heavy metals are lower than 1 except for in Cocoa which is higher than 1 due to the high concentration of Cu and Fe. Therefore, the nuts and seeds analyzed in this study are contaminated with Cu and Fe and so must be taken with care so that consumers will not be exposed to excessive concentration of these metals which may have undesirable effects.

*Corresponding author: E-mail: sirkay006@yahoo.com, drakademola@yahoo.com;

Keywords: Heavy metals; radionuclides; fatal cancer risk; nuts; seeds.

1. INTRODUCTION

The human environment is radioactive and human beings are exposed anywhere in the environment either or both from natural radioactivity, artificial radioactivity, and medical applications. Human beings are exposed from the intake of water, air, soil, and plants [1]. It is a well-known fact that plants used as food commonly contain natural radionuclides and their progenies [2]. Also, other contaminants, like heavy metals are added into plants through the applications of fertilizer, pesticides, herbicides in agriculture, deposition of dust from mining activities, and other industrial activities such as wastes disposal and automobile exhaust [3].

Nuts and seeds are beneficial to humans. It contains many nutrients beneficial to both adults and the young ones. It is consumed as a delicacy and sometimes to treat some diseases and ailments. Medicinal applications of nuts and seeds are limited due to the presence of NORMs and some toxic metals in them [4,5]. Some important information on the level of radionuclides and heavy metals in medicinal plants, nuts, and seeds have been reported through previous researches [6-8]. The process of transfer of radionuclides and heavy metals to the human body is majorly through inhalation and ingestion of contaminated foodstuffs. These contaminants especially radionuclides emit alpha particles that interact with water in the cells and result in some biological effects (mutation). Chronic effects are accompanied by lung cancer, tumor, genetic effects, kidney failure, and vascular diseases [9].

In Nigeria, the side effects of orthodox medicine are changing people's perception about their uses and there is an increase urge in finding other natural means from consumption of natural plants, nuts, and seeds to treat ailments. Therefore, it is imperative to assess the risk that may arise from the consumption of some nuts and seeds to checkmate food fraud [10] and to assess contaminants in them [11]. This study aimed at estimating NORMs, some heavy metals, and evaluation of hazard in the consumption of some nuts and seeds in South-West, Nigeria. The findings of this study would assist the local and international agencies in quality control of food products.

2. MATERIALS AND METHODS

Five samples each of ten nuts and seeds (cashew nut, groundnut, walnut, Kola nut, moringa seed, bitter kola nut, African Star Apple seed, avocado seed, watermelon seed, and cocoa seed) commonly cultivated and consumed in the south-western state of Nigeria were selected for this study. The samples were collected from the farm sites and were packed with identification labels. Nuts and seeds were removed from their shells and were thereafter left to dry at room temperature in the laboratory. The samples were crushed and sieved. 100 g of each sample was weighed and transferred into a cylindrical plastic container and hermetically sealed for thirty days [12] for equilibrium between ^{238}U (^{226}Ra) and ^{232}Th (^{228}Ra) and their respective progenies to take place [13]. The samples were afterward analyzed using the spectrometry method. For heavy metal, their concentration was measured using Inductively Coupled Plasma Spectrometry (ICPS).

2.1 Radionuclide Analysis

The analysis of radionuclide in the samples was achieved using NaI (TI) detector. A Canberra multi-channel analyzer coupled with a scintillating detector was used for the counting. The model no of the detector is 8020 and is manufactured in the USA by Bicron Electronics Ltd. The detector is fixed to a multi-channel analyzer by a coaxial cable. The detector is shielded with a lead shield of about 5 cm thick to prevent radiation from other environmental sources present (background radiation). Gamma sources supplied by the IAEA were used for calibration and the energy-channel calibration was used to fit the linearity graph. A reference source prepared by Rocketdyne Laboratories, Canoga Park, in the USA which is traceable to standard gamma source (No. 48722-356) was used for detection efficiency calibration. The resolution of the detector assembly is about 8% at 0.662 MeV of ^{137}Cs .

The net count was obtained by subtracting the background count from the gross count. The samples were counted for 36000 (10 h). Photopeak energy 1.465 MeV from ^{40}K was used to determine ^{40}K while photopeak energy 1.765 MeV from ^{214}Bi was used to determine ^{226}Ra . ^{232}Th was determined from energy photopeak 2.615 MeV from ^{208}Tl and ^{228}Ra was determined

from gamma-ray energies of its daughter ^{228}Ac (911.07 keV). The concentrations of NORMs in the samples were calculated using a spectrum acquisition and analysis software program (Genie 2K, manufactured by Canberra Industries Inc. USA).

2.2 Heavy Metal Analysis

Samples were digested using the wet digestion method. One gram of each sample was weighed into a 100 mL tube and the tubes were labeled accordingly. Five millilitres of HNO_3 and H_2O_2 were added into the samples and H_2SO_4 was added in drops. The mixture was stirred until the solution becomes clear. The solution was poured out of the tube and mixed with deionized water up to 100 mL and it was then filtered. The toxic metals concentration was obtained with inductively coupled plasma spectrometry (ICPS). Concentrations of metals in the digested samples were determined using plasma and a spectrometer. The solution is directed into the spray chamber through a nebulizer using a peristaltic pump. Each element measured with a spectrometer has its characteristic emission spectrum. The light intensity on the wavelength was measured and with the ICPS calibrated, the concentration of each element was evaluated.

3. RESULTS

3.1 Activity Concentrations of Radionuclides and Calculation of Radiological Parameters

The concentration of the natural radionuclide in nuts and seeds is shown in Table 1. In nuts, the concentration of ^{226}Ra is lowest (3.2 ± 1.0) in the bitter kola nut and highest (10.9 ± 3.2 Bq/kg) in kola nuts. In seeds, it ranged from 4.2 ± 1.6 in African star apple seeds to 14.5 ± 2.8 Bq/kg in Cocoa seeds. It was not detected in groundnut, walnut, moringa seeds, and watermelon seeds. The concentration of ^{228}Ra in nuts is minimum (2.4 ± 0.5 Bq/kg) in Walnut and maximum (5.9 ± 2.4 Bq/kg) in Groundnut. In seeds, the measured activity ranged from 1.8 ± 1.1 in Cocoa seeds to 3.2 ± 1.6 Bq/kg in Avocado seeds and was not detected in bitter kola nut, moringa, and watermelon seeds. The concentration of ^{40}K in nuts ranged from 53.6 ± 4.2 in bitter kola nut to 126.2 ± 22.1 Bq/kg in kola nut. In seeds, it ranged from 50.6 ± 3.5 Bq/kg in Avocado to 123.1 ± 5.6 Bq/kg in African star apple seeds.

3.2 Effective Dose

The annual effective dose from ingestion of nuts and seeds was estimated using the equation

$$E_{Dose} = C_R \cdot DCF_i \cdot A \quad (1)$$

Where DCF_i is dose the conversion factor for ingestion of the three radionuclides analyzed (i.e. 2.8×10^{-7} Sv/Bq, 2.3×10^{-7} Sv/Bq and 6.2×10^{-9} Sv/Bq for ^{228}Ra , ^{226}Ra and ^{40}K respectively for adults), C_R is the consumption rate of NORMs from ingestion of seeds and nuts, the rate from the intake of 1.8 kg/year was assumed [14]. A is the concentration of NORMs in nuts and seeds. The E_{dose} estimated is presented in Table 2, column 1.

3.3 Fatal Lung Cancer Risk and Hereditary Cancer Risk

According to WHO, 2018 [15], Lung cancer causes about 30,000 deaths among non-smokers in the United States alone in 2018. The International Commission on Radiological Protection had proposed several models for calculating lung cancer risk from exposure to radionuclides either through ingestion or inhalation. ICRP proposed 0.05 as a value to calculate risk factors to the public from the ingestion of radionuclides [16]. Fatal Cancer Risk (FCR) from the ingestion of the nuts and seeds was evaluated using equation 2 [17].

$$\text{Fatal Cancer Risk} = ED \times LE \times RF \quad (2)$$

where ED is the mean effective dose ($\mu\text{Sv/y}$), LE is life expectancy (55 years in Nigeria [15] and RF is a factor for risk estimation (Sv^{-1}). The estimated average lifetime fatality cancer risk in nuts and seeds is presented in Table 2, column 2.

Hereditary cancer risk was estimated using the Cancer risk methodology proposed by ICRP [17].

$$\text{Hereditary Cancer Risk} = ED \times LE \times RF \quad (3)$$

where ED is the mean effective dose ($\mu\text{Sv/y}$), LE is the expected life duration (55 years) and RF is a factor for risk estimation (Sv^{-1}). The mean estimated lifetime hereditary cancer risk in the samples is presented in Table 2, column 3.

Table 1. Mean activity concentration (Bqkg⁻¹) of natural radionuclides in seeds and nuts

Samples	⁴⁰ K	²²⁶ Ra	²²⁸ Ra
Kola nut (<i>Cola acoinata</i>)	112	9.2	3.8
Cashew nut (<i>Anacardium occidentale</i>)	111	6.5	2.9
Bitter Kola nut (<i>Garcinia kola</i>)	74	4.0	ND
Groundnut (<i>Arachis hypogeal</i>)	98	ND	5.0
Walnut (<i>Juglans Regia</i>)	96	ND	2.8
Average (Nut)	98.2	6.57	3.63
Avocado seed (<i>Persea americana</i>)	70	9.8	2.8
African Star Apple seed (<i>Chrysophyllum albidum</i>)	113	5.0	2.7
Cocoa seed (<i>Theobroma cacao</i>)	100	10.5	2.2
Moringa seed (<i>Moringa olifera</i>)	102	ND	ND
Watermelon seed (<i>Citrullus lanatus</i>)	93	ND	ND
Average (Seed)	95.6	8.43	2.57

Table 2. Mean annual effective dose (μSvy⁻¹), fatal cancer risk (MPY)⁻¹ and hereditary cancer risk (MPY)⁻¹ in nuts and seeds

Samples	E-dose	Fatal cancer risk	Hereditary cancer risk
Kola nut (<i>Cola acoinata</i>) (<i>Prunusdulcis</i>)	18.05	0.50	0.40
Cashew nut (<i>Anacardium occidentale</i>)	16.36	0.45	0.36
Bitter Kola nut (<i>Garcinia kola</i>)	9.80	0.27	0.21
Groundnut (<i>Arachis hypogeal</i>)	13.30	0.36	0.30
Walnut (<i>Juglans Regia</i>)	11.97	0.33	0.26
Average (Nut)	13.99	0.38	0.30
Avocado seed (<i>Persea americana</i>)	6.25	0.17	0.14
African Star Apple seed (<i>Chrysophyllum albidum</i>)	15.86	0.45	0.35
Cocoa seed (<i>Theobroma cacao</i>)	16.46	0.45	0.36
Moringa seed (<i>Moringa olifera</i>)	11.22	0.31	0.25
Watermelon seed (<i>Citrullus lanatus</i>)	10.23	0.28	0.23
Average (Seed)	12.00	0.33	0.27

3.4 Heavy Metal Analysis

The concentration of some of the heavy metals in the samples is shown in Table 3. Pb and Cd were not detected in a substantial amount or are below the detection limit. The concentration of Cr is only found in the kola nut and not detected in other nuts while the concentration of Iron (Fe) is the highest in all the samples. This might be due to the fact that Fe is an essential element in human life and it bonds with haemoglobin. Cu and Zn concentrations are also higher than recommended in cashew nuts.

In seeds, Fe concentration is very high especially in cocoa and watermelon than the WHO recommendation. High values of Cu, Zn, and Cr were also established in cocoa seeds while Zn concentration is higher than WHO recommendation in African star Apple.

3.5 Calculation of Risks in the Samples

The health risk from the ingestion of heavy metals in the samples (HQ) was estimated as the fraction of quantity of metal taken in a day (ADD in mg/kg of the bodyweight/day) and quantity of metal recommended to be taken per day (RF Do; in mg/kg/day).

$$ADD = \frac{C_{metal} \times I}{D_{average}} \quad (4)$$

where C_{metal} is the concentration of metal; I - daily intake of the metals; $D_{average}$, bodyweight of an adult. The value of ADD estimated for nuts and seeds samples are presented in Table 4. ADD for all metals in the samples is lower than 1

To estimate the health risks, equation 5 was used.

$$HQ = \frac{ADD}{RfD} \quad (5)$$

Where RfD is Oral RfDo. According to USEPA, oral RfD is 1.5 for Cr, 0.04 for Cu, 0.3 for Zn, 0.7 for Fe [18].

If HQ is greater than 1, it means that risk is associated with the intake of the particular metal. The health risk (HQ) obtained from the ingestion of the samples is presented in Table 5. All the HQs obtained in the samples were lower than 1 except for Cu in Cocoa in which HI is higher than 1 indicating a potential risk of exposure to Cu from the consumption of Cocoa.

The Total Hazard Index in the samples was calculated using equation (9) [19].

$$HI = HQ_{Fe} + HQ_{Cu} + HQ_{Zn} + HQ_{Cr} \quad (6)$$

If HI is <1, it means no hazard, if THI is 1.1 – 3, it means probable hazard from consumption of the samples. If HI is 3-10, it is likely to cause a fatal risk [19].

4. DISCUSSION

The concentration of NORMs in this study showed that ^{40}K had the highest contribution in all the samples. The concentration of ^{40}K obtained in this study is within the world average. The concentration of ^{226}Ra was highest in kola nut but below the detection limit in groundnut and walnut. The concentration of ^{228}Ra was also below the detection limit in bitter kola nut. The mean concentrations of the radionuclide analyzed in nuts were respectively, 98.3 ± 13.5 , 6.6 ± 1.8 , and 3.6 ± 1.0 Bq/kg for ^{40}K , ^{226}Ra , and ^{228}Ra . These values are within the recommended values of NORMs in food and food products [1].

Table 3. Mean concentration of heavy metals in nuts and seeds


Samples	Fe (mgkg ⁻¹)	Cu (mgkg ⁻¹)	Zn (mgkg ⁻¹)	Cr (mgkg ⁻¹)
Nuts				
Kola nut (<i>Cola acominata</i>)	9.28	9.28	27.67	0.08
Cashew nut (<i>Anacardium occidentale</i>)	26.95	26.95	50.40	ND
Bitter Kola nut (<i>Garcinia kola</i>)	18.68	3.07	2.75	ND
Groundnut (<i>Arachis hypogea</i>)	17.62	8.15	23.98	ND
Walnut (<i>Juglans regia</i>)	28.86	15.99	37.22	ND
Seeds				
Avocado seed (<i>Persea americana</i>)	93.45	10.58	14.22	0.52
African Star Apple seed (<i>Chrysophyllum albidum</i>)	63.46	13.82	46.37	0.36
Cocoa seed (<i>Theobroma cacao</i>)	542.49	51.57	65.82	5.11
Moringa seed (<i>Moringa olifera</i>)	51.96	3.96	32.29	0.08
Watermelon seed (<i>Citrullus lanatus</i>)	327.31	7.42	34.35	0.29

Table 4. Average Daily Dose (ADD) of metals in the nuts and seeds samples

Samples	Fe(mgkg ⁻¹) x 10 ⁻³	Cu(mgkg ⁻¹) x 10 ⁻³	Zn(mgkg ⁻¹) x 10 ⁻³	Cr(mgkg ⁻¹) x 10 ⁻⁴
Nuts				
Kola nut (<i>Cola acominata</i>)	8.62	8.62	25.70	0.74
Cashew nut (<i>Anacardium occidentale</i>)	25.1	25.1	47.80	ND
Bitter Kola nut (<i>Garcinia kola</i>)	17.40	2.85	2.55	ND
Groundnut (<i>Arachis hypogea</i>)	16.36	7.57	22.27	ND
Walnut (<i>Juglans regia</i>)	26.79	14.85	34.56	ND
Seeds				
Avocado seed (<i>Persea americana</i>)	86.77	9.82	13.20	4.83
African Star Apple seed (<i>Chrysophyllum albidum</i>)	58.92	12.83	43.06	3.34
Cocoa seed (<i>Theobroma cacao</i>)	503.40	47.89	61.12	47.45
Moringa seed (<i>Moringa olifera</i>)	14.80	0.37	0.30	0.74
Watermelon seed (<i>Citrullus lanatus</i>)	303.93	6.89	31.90	2.69

Table 5. Hazard quotient (HQ, (mg/kg)) and fractional hazard index (HI) of heavy metals in nuts and seeds samples

Samples	Fe (mgkg ⁻¹)	Cu (mgkg ⁻¹)	Zn (mgkg ⁻¹)	Cr (mgkg ⁻¹)x 10 ⁻⁴	HI = \sum HQ
Nuts					
Kola nut (<i>Cola acominata</i>)	0.012	0.216	0.086	0.495	0.216
Cashew nut (<i>Anacardium occidentale</i>)	0.036	0.626	0.156	ND	0.818
Bitter Kola nut (<i>Garcinia kola</i>)	0.025	0.071	0.009	ND	0.105
Groundnut (<i>Arachis hypogeal</i>)	0.023	0.189	0.074	ND	0.286
Walnut (<i>Juglans regia</i>)	0.038	0.371	0.115	ND	0.524
Seeds					
Avocado seed (<i>Persea americana</i>)	0.124	0.246	0.044	3.219	0.414
African Star Apple seed (<i>Chrysophyllum albidum</i>)	0.084	0.321	0.144	2.229	0.549
Cocoa seed (<i>Theobroma cacao</i>)	0.720	1.197	0.204	31.633	2.121
Moringa seed (<i>Moringa olifera</i>)	0.069	0.092	0.099	0.495	0.260
Watermelon seed (<i>Citrullus lanatus</i>)	0.434	0.172	0.106	1.795	0.712

**Fig. 1. Percentage concentrations of radionuclide in nut samples**

In seeds, the concentration of ⁴⁰K was highest in African Star Apple and lowest in Avocado with a mean of 95.7±15.0 Bq/kg. ²²⁶Ra concentration was below the detection limit in Moringa and watermelon seeds. The maximum concentration of ²²⁶Ra was from Cocoa seeds. The mean concentration of ²²⁶Ra in seeds was obtained as 8.4±2.6 Bq/kg. The concentration of ²²⁸Ra in the seeds was found to be lower than the recommendation in all the samples with the highest from avocado. The concentration of ⁴⁰K, ²²⁶Ra, and ²²⁸Ra obtained in this study are within the recommended average [1]. The

concentration of NORMs in the samples follow the order ⁴⁰K > ²²⁶Ra > ²²⁸Ra as shown in Fig. 1 and 2. The percentage concentrations of ²²⁶Ra, ²²⁸Ra and ⁴⁰K in nuts and seeds are the same and so there is no significant difference in the concentration of radionuclides in nuts and seeds as shown in Figs. 1 and 2. The mean activity of radionuclides measured was compared using a 2-tail paired t-test at a 95% level of significance (p-value = 1.645) and it was observed that there is no significant difference between their concentration (Calculated value = 0.29, -1.28 and 1.36, respectively).




Fig. 2. Percentage concentrations of radionuclide in seed samples

Effective dose estimated from the ingestion of nuts and seeds confirmed that the largest effective dose was from kola nuts ($18.05 \mu\text{Svy}^{-1}$) while the lowest effective dose was from Avocado ($6.25 \mu\text{Svy}^{-1}$). The mean annual effective doses for nuts and seeds are 13.99 and $12.00 \mu\text{Svy}^{-1}$, respectively (Table 2). The average value obtained in this study is within the limit recommended by ICRP [17].

The Fatal Lung Cancer Risk and hereditary cancer risk estimated from the ingestion of nuts and seeds in this study showed that an average of 0.38 and 0.30 people in a million will contract cancer from consumption of nuts and 0.33 and 0.27 people will contract cancer from the consumption of seeds. The highest risk estimated was from consumption of kola nuts while the lowest risk was from consumption of bitter kola nuts. The values obtained in this study are below the world average of one person in a million per year [18,20].

The analyses of metals in the samples revealed that Cr had the lowest in all samples and not detectable in Cashew nut, Bitter kola nuts, Groundnut, and Walnut. The concentrations of Chromium obtained in these samples are lower than the regulation [21]. In Nuts, The concentration of Fe was lowest in Groundnut (17.62 mg/kg) and highest in Walnut (28.86 mg/kg) and in seeds, Fe concentration was highest in Cocoa (542.49 mg/kg) and lowest in

Moringa seed. It was also observed that the concentration of Fe is higher in Cocoa and Mustard seeds than the World Health Organization recommendations [21]. Copper (Cu) concentration in the samples ranged from 3.07 mg/kg in Bitter kola nuts to 26.95 mg/kg in Cashew nut. The concentration of Cu was found to be higher than 10 mg/kg recommendation in Cashew and Walnuts [22]. Also in seeds, Cu was found to be higher than the recommended concentration in Avocado, African star Apple and Cocoa seed [22]. The concentration of Zinc (Zn) ranged from 2.75 mg/kg in Bitter kola nuts to 50.4 mg/kg in Cashew nuts. Zn concentration was also found to be higher than 40 mg/kg which is the upper intake for a healthy adult in Cashew, African star Apple and Cocoa seeds [20,23]. The concentrations of Cu, Cr, and Zn were also observed to be higher than the WHO recommendations in Cocoa [21].

The Hazard Quotient (HQ) and Hazard Index (HI) estimated for all samples are lower than 1 except for Cocoa in which HI exceeded 1. This implies that there may be a hazard in Cocoa seeds consumption due to the high concentration of Cu. Although, Copper is a vital dietary nutrient in which only small amounts are needed for human well-being. In high doses, Cu can be extremely toxic [24]. A high dose of Zn may also lead to tachycardia, vomiting, diarrhea, pancreatitis, and damage of hepatic parenchyma [25].

5. CONCLUSION

Measurement of natural radionuclides and some heavy metals in some nuts and seeds commonly consumed in south-western parts of Nigeria was carried out in this study. The Radioactive analysis was carried out using NaI (TI) detector. The results of the analysis showed that the concentrations of ^{226}Ra , ^{228}Ra , and ^{40}K were within the recommended average. The estimated effective doses were found to be lower than the recommended values. The assessment of the cancer risks from the consumption of nuts and seeds showed that the risks estimated are lower than the safe limit of 1×10^{-6} to 1×10^{-4} [18,20]. Therefore, no radioactive health risk was involved in the consumption of nuts and seeds if consumed within the 42 g per day recommended.

The concentration of heavy metals was measured using Inductively Coupled Plasma (ICP). The concentration of heavy metals in the samples followed the order $\text{Fe} > \text{Zn} > \text{Cu} > \text{Cr}$. Concentration of Zn was found to be higher than the recommendation in Cashew, African star Apple and Cocoa seeds. The measured Cu concentration was found to be higher than the recommended value in Cashew nuts and Walnuts. The Hazard Indices estimated from consumption of heavy metals in the samples are lower than 1 except for in Cocoa where HI is higher than 1. Therefore, nuts and seeds analyzed in this study are contaminated with some heavy metals and so must be taken with care because excessive intake of iron may increase susceptibility to infection; excessive exposure to copper might result in liver cirrhosis and kidney damage [26] while high dose of Zn may cause tachycardia, vomiting, diarrhea, pancreatitis, and damage of hepatic parenchyma [25].

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. United Nations Scientific Committee on the effects of atomic radiation: Sources and effects of ionizing radiation United Nations Scientific Committee on the Effects of Atomic Radiation. United Nations, New York. UNSCEAR; 2000. Available:www.unscear.org
2. International food safety authorities network. Information on nuclear accidents and radioactive contamination of foods international food safety authorities network. World Health Organization (WHO), Geneva. INFOSAN; 2011. Available:www.fao.org/3/a-i3345e.pdf
3. Jarup L. Hazards of heavy metals contamination, *Bri. Med. Bull.* 2003;68(1): 167-82. Available:https://doi.org/10.1093/bmb/ldg032
4. Mahurpawar M. Effects of heavy metals on human health, *Int. J. Res. Granthaalayah.* 2015;1-7.
5. Lanre-Iyanda TY, Adekunle IM, Assessment of heavy metals and their estimated daily intakes from two commonly consumed foods (Kulikuli and Robo) found in Nigeria. *Africa J. Food. Agri. Nutr. Devt.* 2012;12(3):6156-6169. eISSN: 1684-5374
6. Ademola AK, Omoboyede JO Health risk assessment of natural radionuclide and heavy metals in commonly consumed medicinal plants in South-West Nigeria. *Ife J. Sci.* 2018;20(3):529-537. Available:https:// DOI:10.4314/ijfs.v20i3.7
7. Yin LL, Tian Q, Shaq XZ, Kong XY, Qiu JY. Determination of trace elements in edible nuts in the beijing market by ICP-MS. *Biomed. Envir Sci.* 2015;28(6):449-454. DOI:https://DOI: 10.3967/bes2015.063
8. Chung KH, Shin KO, Hwang HJ, Choi KS, Chemical composition of nuts and seeds sold in Korea. *Nat. Res Prac.* 2013;7(2), 82– 88. DOI:https://doi.org/10.4162/nrp.2013.7.2.82
9. El-Zakla T, Abdel-Ghny HA, Hassan AM. Natural radioactivity of some local fertilisers. *Romanian J. Phys.* 2007;52: 731–739
10. Johnson R. Food fraud and economic motivated adulteration of food and food ingredients, Congressional Research Service of US Department of Agriculture. Congressional Research Service. 2014;7-5700. Available:https://fas.org/spp/crs/misc/R43358.pdf
11. European Food Safety Authority. Cadmium. Scientific opinion of the panel on contaminants in the food chain of the European Food chain (CONTAM Panel). EFSA. 2009;J.980:1-139.

- DOI: 10.2903/j.efsa.2009.980
Available: <https://www.efsa.europa.eu/en/efsajournal/pub/980>
12. Atomic Energy Regulatory Board. Accreditation of laboratory for measurement of radionuclide content in commodities. Atomic Energy Regulatory Board, Mumbai; 2003.
Available: <https://www.aerb.gov.in>
 13. Sathyapriva RS, Rao DD, Prabhath RK. Choosing an appropriate method for measurement of ²³²Th in environmental samples. *Radiat. Prot. Environ.* 2017;40: 90-94.
DOI: https://doi.org/10.4103/rpe.RPE_20_17
 14. Harb SRM. Natural radioactivity concentrations in some medicinal plants and annual committed effective dose from their consumption. *Radiat. Prot. Environ.* 2015;38(1&2):35-38.
Available: <https://doi.org/10.4103/0972-0464.162816>
 15. World Health Organisation (WHO) Data published for Life Expectancy in Nigeria; 2018.
Available: www.who.int/countries/nga
 16. International Commission on Radiological Protection. 1990 Recommendation of the International Commission on radiological protection. Oxford: Pergamon Press. ICRP publication 60, *Annals of the ICRP*. 1991; 21(1-3).
Available: www.icrp.org/docs/Histpol.pdf
 17. Taskin H, Karavus M, Ayb B, Topuzoglu A, Hidiroglu S, Karahan G. Radionuclide concentrations in soil and lifetime cancer risk due to gamma radioactivity in Kirklareli, Turkey. *J. Environ. Radioactiv.* 2009;100:49–53.
Available: <https://doi.org/10.1016/j.jenvrad.2008.10.012>
 18. United State Environmental Protection Authority, Diffuse NORM waste characterization and preliminary risk assessment. S. Cohen and Associates, Inc. Rogers & Associate Engineering Corp. US Environmental Protection Agency. Office of Radiation and Indoor Air (ANR-459);1993.
Available: <https://nepis.epa.gov>
 19. United States Environmental Protection Agency. EPA regional screening level (RSL) summary table. Washington, DC; 2011.
Available: <https://www.epa.gov/risk/regional-screening-levels>
 20. World Health Organization. Guidelines on assessing quality of herbal medicines with reference to contaminants and residues. Geneva, World Health Organization; 2007.
Available: apps.who.int/iris/handle/10665/43510
 21. Food and Agriculture Organization/World Health Organization (FAO/WHO). Joint Expert Committee on Food Additives. Summary and conclusions. 53rd Meeting, Rome; 1999.
Available: www.fao.org/fileadmin/templates
 22. Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological profile for Cu. U.S. Department of Health and Human Services. Public Health Service; 2004.
Available: www.atsdr.cdc.gov/toxprofiles
 23. Muhammad A, Sreebas CB. Assessment of heavy metals concentration in some selected medicinal plants collected from BCSIR, Chittagong Cultivation Area in Bangladesh, *Hamdard Medicus*. 2012; 55(3):26-32.
ISSN: 0250-7188
 24. Ashish B, Necti K, Himanshu K. Copper toxicity: A comprehensive study. *Research J. Recent Sci.* 2013;2:58–67.
DOI: <https://doi.org/10.1360/972012-1626>
 25. Adefila EO, Onwordi CT, Ogunwande IA. Level of heavy metals uptake on vegetables planted on poultry droppings dumpsite, *Arch. Appl. Sci. Res.* 2010;2(1): 347-353.
 26. Gaetke LM, Chow CK. Copper toxicity, oxidative stress, and antioxidant nutrients. *Toxicology*. 2003;189:147–163.
DOI: [https://doi.org/10.1016/S0300-483X\(03\)00159-8](https://doi.org/10.1016/S0300-483X(03)00159-8)

© 2020 Ademola and Morakinyo; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/58704>